JC07 Rec'd PCT/PTO 0 9 JAN 2002

ATTORNEY'S DOCKET NUMBER FORM PTO-1390 U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE (MODIFIED) G-1344 U.S. APPLICATION NO. (if known, see 37 C.F.R. 1.5) TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) 10/031122 **CONCERNING A FILING UNDER 35 U.S.C. 371** INTERNATIONAL APPLICATION NO. INTERNATIONAL FILING PRIORITY DATE CLAIMED DATE PCT/GB00/02361 06/15/2000 (06.15.00) 07/13/1999 (07.13.99) TITLE OF INVENTION: SULFONAMIDE SUBSTITUTED BENZYLAMINE DERIVATIVES AND THEIR USE AS **MEDICAMENTS** APPLICANT(S) FOR DO/EO/US: Sandra Ginette Milutinovic, et al. Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information: This is a FIRST submission of items concerning a filing under 35 U.S.C. 371. 1. This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371. 2. 3. This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1). A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date. A copy of the International Application as filed (35 U.S.C. 371(c)(2)) Ļ is transmitted herewith (required only if not transmitted by the International Bureau). has been transmitted by the International Bureau. b. fĻ is not required, as the application was filed in the United States Receiving Office (RO/US). Ŋ A translation of the International Application into English (35 U.S.C. 371(c)(2)). Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3)) are transmitted herewith (required only if not transmitted by the International Bureau). a. have been transmitted by the International Bureau. b. have not been made; however, the time limit for making such amendments has NOT expired. c. M d. have not been made and will not be made. 8. A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)). 9. X An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). A copy of the International Preliminary Examination Report (IPER), including any annexes, and, if not in English, an 10. X English language translation of the annexes to the IPER under PCT Article 36 (35 U.S.C. 371(c)(5)). Items 11. to 16. below concern document(s) or information included: An Information Disclosure Statement under 37 CFR 1.97 and 1.98. 12. An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. 13. A FIRST preliminary amendment. A SECOND or SUBSEQUENT preliminary amendment. 14. A substitute specification. 15. A change of power of attorney and/or address letter. 16. Other items or information:

JC13 Rec'd PCT/PTO 0 9 JAN 2002

U.S. APPLICATION NO. (i	f known, see 37 C.F.R. 1.5)	ATTORNEY'S DOCKET NUMBER G-1344				
17. X The following	ng fees are submitted:			CALCULATIONS	PTO USE ONLY	
Neither international	NAL FEE (37 CFR 1.49 onal preliminary examina search fee (37 CFR 1.44 Search Report not prepa	ation fee (37 CFR 1.482 5(a)(2)) paid to USPT(Ó			
	liminary examination fee Search Report prepared					
	liminary examination feerch fee (37 CFR 1.445(a)					
	liminary examination fee I not satisfy provisions of					
	liminary examination fee aims satisfied provisions					
in the second se	ENTED ADDDOD		E AMOUNT -	\$ 890.00	Ι	
Surcharge of \$130.00	ENTER APPROPI			\$ 890.00	 	
months from the earlie	st claimed priority date ((37 CFR 1.492(e)).		φ 	<u> </u>	
CLAIMS Total claims	NUMBER FILED	NUMBER EXTRA	RATE	1 1 2 6 0 0	,	
1.5	27 - 20=	7	X \$18.00	\$ 126.00		
ndependent claims	2 - 3=		X \$84.00	\$		
A CONTRACTOR OF THE CONTRACTOR	DENT CLAIM(S) (if app	OF ABOVE CAL	+ \$280.00	\$ 280.00	<u> </u>	
	\$					
	iling by small entity, if a e filed (Note 37 CFR 1.9			\$		
			SUBTOTAL =	\$ 1296.00		
	0.00 for furnishing Englished claimed priority date ((37 CFR 1.492(f)).	+	\$		
			IONAL FEE =	\$		
	enclosed assignment (37 propriate cover sheet (37	7 CFR 3.28, 3.31).	ignment must be 40.00 per property	\$		
+			40.00 per property			
		TOTAL FEES	ENCLOSED =	\$ 1296.00		
				Amount to be	\$	
				refunded	<u> </u>	
A =1-==1- 1: 6	ha amount of the	40 0000m 411 C		charged	\$	
b. X Please charg	he amount of \$ ge my Deposit Account N sheet is enclosed.	_ to cover the above for the open to the amount of the amo		cover the above fees.	A duplicate	
c. X The Commis	ssioner is hereby authorize t to Deposit Account No				t any	
	opropriate time limit ur 37(a) or (b)) must be fil				ive	
SEND ALL CORRESPO ELI LILLY AND COM PATENT DIVISION/JA	IPANY G	<u> </u>	Jant 9. x SIGNATI	Gongola URE		
LILLY CORPORATE C	ENTER	 -	Janet A. G NAME	,		
Date		D E	48,436 GISTRATION NUMB	(317) 277-0 FR TELEPHON	6186 JE NUMBER	

Rec'd PCT/PTO 0 9 JAN 2002

SULFONAMIDE SUBSTITUTED BENZYLAMINE DERIVATIVES AND THEIR USE AS MEDICAMENTS

This invention relates to novel chemical compounds and their use as pharmaceuticals.

5

It is well known that chemical compounds which modulate the activity of neuronal calcium channels are potentially useful in treating disorders of the central nervous system.

10

The compounds of the invention have the following general formula:

20

in which the aminosulfonyl group is attached at the 3or 4-position, and in which

 R^1 is hydrogen, C_{1-6} alkyl, C_{3-10} cycloalkyl, C_{3-10} cycloalkyl- C_{1-4} alkyl or optionally substituted phenyl- C_{1-4} alkyl,

 $\rm R^2$ is $\rm C_{1-6}$ alkyl, $\rm C_{3-10}$ cycloalkyl, $\rm C_{3-10}$ cycloalkyl- $\rm C_{1-4}$ alkyl, optionally substituted phenyl- $\rm C_{1-4}$ alkyl or -(CH₂) $\rm _2NR^5R^6$ where $\rm R^5$ and $\rm R^6$ are each hydrogen or $\rm C_{1-6}$ alkyl, and

 R^3 and R^4 are each C_{1-6} alkyl, C_{3-10} cycloalkyl- C_{1-4} alkyl, C_{3-6} alkenyl, optionally substituted phenyl or optionally substituted phenyl- C_{1-4} alkyl,

or \mathbb{R}^1 and \mathbb{R}^2 , or \mathbb{R}^3 and \mathbb{R}^4 , or \mathbb{R}^5 and \mathbb{R}^6 , together with the nitrogen atom to which they are attached, form a carbocyclic group containing 4 to 7 carbon atoms optionally substituted with one to three methyl or ethyl groups and optionally containing an oxygen atom or a further nitrogen atom, said carbocyclic group being

optionally fused to an optionally substituted phenyl group;

or a salt thereof.

5

The compounds of the invention have been found to be active in tests that show modulation of voltage-dependent calcium channels, and are thus indicated for use in the treatment of diseases in which such modulation is beneficial, in particular disorders of the central nervous system.

Thus, the invention includes a method of treating a disorder of the central nervous system, which comprises administering an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof.

The invention also includes the use of a compound of

formula (I), or a pharmaceutically acceptable salt

thereof, in the manufacture of a medicament for treating
a disorder of the central nervous system.

The invention also includes the use of a compound of formula (I), or a pharmaceutically acceptable salt

thereof, in the manufacture of a medicament for treating a disorder of the central nervous system.

In the above formula (I), a C_{1-6} alkyl group includes methyl, ethyl, propyl, isopropyl, butyl, tert. butyl and hexyl, and is preferably methyl or ethyl. A substituted phenyl group is phenyl substituted with one or more, for example one to three, substituents selected from, for example C_{1-4} alkyl, especially methyl, C_{1-4} alkoxy,

especially methoxy and ethoxy, hydroxy, nitro, cyano, halo, especially chloro or fluoro, trihalomethyl, especially trifluoromethyl, carboxy and C_{1-4} alkoxycarbonyl. A halo atom is preferably chlorine, bromine or fluorine. A substituted phenyl group preferably has one to three substituents selected from hydroxy, C_{1-4} alkyl, halo, nitro and trifluoromethyl. An optionally substituted phenyl- C_{1-4} alkyl group is preferably of the formula $R-(CH_2)_n-$ where R is optionally substituted

phenyl and n is 1 to 4, but the linking chain can also

- be branched alkylene. A C_{3-10} cycloalkyl group is preferably, for example, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl and these groups may optionally be substituted by one or two C_{1-4} alkyl, especially methyl, substituents. A C_{3-10} cycloalkyl-
- 25 C_{1-4} alkyl group is one such cycloalkyl group attached

to a C_{1-4} alkyl, and is preferably of the formula $R-(CH_2)_n-\mbox{ where R is cycloalkyl and n is 1 to 4.} \mbox{ When }$ or R^4 is C_{1-6} alkyl it is preferably C_{3-6} alkyl.

The groups R¹ and R², R³ and R⁴, and R⁵ and R⁶, can form a carbocyclic ring with the nitrogen to which they are attached and optionally also contain an oxygen atom or an additional nitrogen. Preferred examples, including the nitrogen of the amino sulfonyl group, are pyrrolidino, piperazino, morpholino and especially 3,5-dimethylpiperidino.

A particular group of compounds of the invention is one of formula (I) in which R^1 , R^2 , R^3 and R^4 are each C_{1-6} alkyl, C_{3-10} cycloalkyl, C_{3-10} cycloalkyl- C_{1-4} alkyl or optionally substituted phenyl- C_{1-4} alkyl, and R^1 can in addition be hydrogen, or R^1 and R^2 , or R^3 and R^4 together with the nitrogen atom to which they are attached, form a carbocyclic group as defined above.

20

15

In a preferred group of compounds R^1 , R^2 , R^3 and R^4 are each C_{1-6} alkyl, C_{3-10} cycloalkyl, C_{3-10} cycloalkyl- C_{1-4} alkyl or optionally substituted phenyl- C_{1-4} alkyl, and R^1 is in addition hydrogen.

It is preferred that R^1 is hydrogen. Furthermore, R^3 and R^4 , which can be the same or different, are preferably C_{1-4} alkyl. It is further preferred that R^2 is optionally substituted phenyl- C_{1-4} alkyl.

A further preferred group of compounds is one of formula (I) in which R^2 is $-(CH_2)_2NR^5R^6$.

A further preferred group of compounds is one of formula (I) in which R³ or R⁴ is C₃₋₆ alkyl or when R³ and R⁴ are taken together with the nitrogen atom they form a piperidine ring which is substituted at the 3-and/or 5-positions with one or two methyl or ethyl substituents.

It will be appreciated that the compounds of the invention can contain one or more asymmetric carbon atom which gives rise to enantiomers. The compounds can be prepared as racemates or can be made from enantiomeric intermediates. Both racemates and enantiomers form part of the present invention.

It will also be understood that salts of the compounds
of the invention can be prepared and such salts are

10

15

WO 01/04087

included in the invention. They can be any of the well known acid addition salts. Acid addition salts are preferably the pharmaceutically acceptable non-toxic addition salts with suitable acids, such as those with inorganic acids, for example hydrochloric, hydrobromic, nitric, sulfuric or phosphoric acids, or with organic acids, such as organic carboxylic acids, for example glycollic, maleic, fumaric, malic,oxalic, tartaric, citric, salicylic or o-acetoxybenzoic acids, or organic sulfonic acids, methane sulfonic, 2-hydroxyethane sulfonic, toluene-p-sulfonic or naphthalene-2-sulfonic acids.

In addition to pharmaceutically-acceptable salts, other salts are included in the invention. They may serve as intermediates in the purification of compounds or in the preparation of other, for example pharmaceutically-acceptable, salts, or are useful for identification, characterisation or purification.

20

The invention includes a process for producing the compounds of formula (I) above which comprises reducing a compound of the fomula:

10

The reaction is preferably carried out in an organic solvent, for example, at a temperature of 0° C. to 100° C., employing a reducing agent, for example lithium aluminium hydride.

Compounds of formula (II) can readily be prepared by conventional methods, for example, by reacting a compound of the formula:

where X is a leaving group such as, for example, halo or hydroxy, with an amine of the formula HNR^1R^2 .

The reaction is preferably carried out in an organic solvent such as, for example, chloroform or acetonitrile, at a temperature of from 0° C. to 100° C. such as, for example, ambient temperature.

10

15

5

Intermediate compounds of formula (III) are known in the art and can be readily prepared by known methods. When an acid halide is employed (X is halo such as, for example, chloro), the reaction is preferably carried out in the presence of a solid phase scavenger to absorb the acid liberated by the reaction. When the free acid is employed (X is hydroxy), a condensing reagent such as,

for example, dimethylaminopropyl-ethylcarbodiimide can be employed.

A further route to the compounds of the invention, which
is also included in the invention, involves the
reduction of the imine corresponding to the compound of
formula (III):

10

15

employing a reducing agent as, for example, sodium borohydride. Compounds of formula (IV) can readily be prepared by reacting an amine of formula R^1R^2NH with the appropriate benzaldehyde derivative, which can, in its turn, be prepared by reducing the corresponding benzoic

acid derivative to the alcohol, followed by oxidation to the required benzaldehyde intermediate.

Amine reactants of the formula $\mathrm{HNR}^1\mathrm{R}^2$ are well known and can be readily prepared by known methods. Those in which R^2 is $-(\mathrm{CH}_2)_2\mathrm{NR}^5\mathrm{R}^6$ can, for example, be prepared by reductive amination, that is, by reacting the appropriate diamine with an aldehyde in reducing conditions.

10

Alternatively, compounds of formula (I) in which R^2 is $-(CH_2)_2NR^5R^6$ can be prepared by alkylation of the corresponding compound of formula (I) in which R^1 is hydrogen.

15

As mentioned above, the compounds of the invention are active in tests that indicate their utility in the treatment of diseases of the central nervous system.

The compounds modulate the activity of calcium channels and, in particular, they block voltage sensitive calcium channels as determined in a test based on Boot J. R., et al., Specificity of autoantibodies in the Lambert-Eaton Myasthenic Syndrome, Ann NY Acad. Sci. (1997), in which measurements of calcium flux using calcium

III THE THE THE THE THE THE

5

10

15

20

WO 01/04087 PCT/GB00/02361
- 12 -

sensitive dyes are made. Compounds described in the following Examples were found to inhibit voltage-dependent calcium channels in cloned human cell lines expressing specific voltage-dependent calcium channels with an IC_{50} of less than 10 μM .

The compounds of the invention are thus indicated for use in the treatment of anoxia, ischaemia, stroke and heart failure, migraine, diabetes, cognitive impairment, pain, epilepsy, traumatic head or spinal injury, AIDS related dementia and blindness, amnesia, neurodegenerative diseases such as Alzheimer's, Parkinson's and Huntington's diseases and age-related memory disorders, Down's syndrome, mood disorders, drug or alcohol addition withdrawal, nausea from chemotherapy, and carbon monoxide or cyanide poisoning.

The invention also includes a pharmaceutical composition comprising a pharmaceutically acceptable diluent or carrier in association with the compound of the invention or a pharmaceutically acceptable salt or ester thereof.

The compound may be administered by various routes, for example by the oral or rectal route, topically or parenterally, for example by injection or infusion,

10

WO 01/04087 PCT/GB00/02361
- 13 -

being usually employed in the form of a pharmaceutical composition. Such compositions are prepared in a manner well known in the pharmaceutical art and comprise at least one active compound. In making the compositions of the present invention, the active ingredient will usually be mixed with a carrier, or diluted by a carrier, and/or enclosed within a carrier which may, for example, be in the form of a capsule, sachet, paper or other container. When the carrier serves as a diluent, it may be a solid, semi-solid, or liquid material which acts as a vehicle, excipient or medium for the active ingredient. Thus, the composition may be in the form of tablets, lozenges, sachets, cachets, elixirs, suspensions, ointments containing, for example, up to 10% by weight of the compound, soft and hard gelatin capsules, suppositories, injection solutions and

Some examples of suitable carriers are lactose,

dextrose, sucrose, sorbitol, mannitol, starches, gum

acacia, calcium phosphate, alginates, tragacanth,

gelatin, syrup, methyl cellulose, methyl- and propyl
hydrobenzoate, talc magnesium stearate and mineral oil.

The compositions of the injection may, as is well known

in the art, be formulated so as to provide quick,

suspensions and sterile packaged powders.

20

5

WO 01/04087 PCT/GB00/02361

sustained or delayed release of the active ingredient after administration to the patient.

Where the compositions are formulated in unit dosage form, it is preferred that each unit dosage form contains from 5 mg to 500 mg. The term 'unit dosage form' refers to physically discrete units suitable as unit dosages for human subjects and animals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with the required pharmaceutical carrier.

The active compound is effective over a wide dosage range and, for example, dosages per day will normally fall within the range of from 0.5 to 300 mg/kg, more usually in the range of from 5 to 100 mg/kg. However, it will be understood that the amount administered will be determined by the physician in the light of the relevant circumstances including the conditions to be treated, the choice of compound to be administered and the chosen route of administration, and therefore the above dosage ranges are not intended to limit the scope of the invention in any way.

The invention is illustrated by the following Preparations and Examples.

EXAMPLE 1

5 4-[(N,N-di-n-propylamino)sulfonyl]-benzoic acid

To a stirred solution of di-n-propylamine (3.03 g, 0.03 mole) in dry tetrahydrofuran (20 ml) at 0° C. (ice/salt bath), was added 4-chlorosulfonylbenzoic acid 10 (2.2 g, 0.01 mole). Stirring was continued for 1 hour. Ice water was added cautiously and the reaction made acid with 2NHCl. The 4-[(N,N-di-n-propylamino)sulfonyl]-benzoic acid was collected by filtration as a white solid which was dried in vacuo at 40° C.

EXAMPLE 2

4-[(N-di-n-propylamino)sulfonyl]-N-4-methoxybenzylbenzamide

To a solution of 4-[(N,N-di-n-propylamino)sulfonyl]benzoic acid (2.85 g, 0.01 mole) in dry dichloromethane

(ml) at 0° C. was added oxalyl chloride (2.54 g,

0.02 mole) and dimethylformamide (4 drops). The

reaction mixture was stirred for 2 hours. The reaction was evaporated to dryness in vacuo. The resulting acid chloride was added to a stirred solution of p-methoxybenzylamine (1.51 g, 0.011 mole) and triethylamine (1.11 g, 0.011 mole) in dry tetrahydrofuran (25 ml) at 0-5° C. After stirring for 4 hours the reaction was poured into ice water and extracted with ethyl acetate. The solvent was washed with brine, dried and evaporated to dryness in vacuo. Chromatography on flash silica using 10% ethyl

10 Chromatography on flash silica using 10% ethyl acetate/dichloromethane gave 4-{(N,N-di-n-propylamino)sulfonyl]-N-4-methoxybenzyl-benzamide as a white solid. M.p. 132-134° C.

15

EXAMPLE 3

N, N-di-n-propyl-4-{[(4-methoxybenzyl)amino]methyl}
benzenesulfonamide

20

25

To a stirred solution of 4-[(N,N-di-n-propylamino)sulfonyl]-N-4-methoxybenzyl-benzamide(1.87g, 4.62mmole) in dry ether (50ml) was added a solution of 2M lithium aluminium hydride in tetrahydrofuran (4.63ml, 9.24mmole). The reaction was heated at reflux for 2 hours. After cooling to room temperature water (1ml)

was added dropwise with caution followed by 2NNaOH

(lml). When gas evolution ceased the reaction mixture

was filtered through a pad of celite which was well

washed with ether. After removal of the solvent in
vacuo the product was purified by chromatography on

flash silica eluting with 10% methanol/ethyl acetate.

The resulting amine was converted to the maleic acid

salt and re-crystallised from ethanol/ether to give N,N
di-n-propyl-4-{[(4-methoxybenzyl)amino]methyl}

10 benzenesulfonamide maleate. mp. 133-135°C

Similarly prepared were:

N, N-di-n-propyl-3-{[(4-methoxybenzyl)amino]methyl}

- benzenesulfonamide maleate. mp. 160-162°C

 N,N-di-n-propyl-4-{[(3,4-dimethoxyphenethyl)]

 amino]methyl}benzenesulfonamide maleate. mp. 130-132°C

 N-(3,3-dimethylpiperidino)-3-{[(4-fluorobenzyl)]

 amino]methyl}benzenesulfonamide maleate. mp. 169-171°C
- N-(3,3-dimethylpiperidino)-4-{[(4-fluorobenzyl)]
 amino]methyl}benzenesulfonamide maleate. mp. 196-198°C
 N,N-di-n-propyl-3-{[(4-fluorobenzyl)amino]methyl}
 benzenesulfonamide maleate. mp. 168-170°C
 N-phenyl-N-n-propyl-4-{[dimethylamino]methyl}
- 25 benzenesulfonamide maleate. mp. 154-156 $^{\circ}$ C

```
N-(3,3-dimethylpiperidino)-3-{[(4-fluorobenzyl)-N-
    methylamino]methyl}benzenesulfonamide maleate.
    spectrum:MH+=405 (TSP+)
    N-(3,3-dimethylpiperidino)-3-{[(4-fluorobenzyl)-N-
    benzylamino]methyl}benzenesulfonamide maleate. mp. 183-
 5
    185°C
    N-phenyl-N-methyl-3-{[(4-fluorobenzyl)amino]methyl}
    benzenesulfonamide maleate. mp. 194-196°C
    N-phenyl-N-n-butyl-4-{[hexylamino]methyl}
    benzenesulfonamide maleate. mp. 106-108^{\circ}C
10
    N-(3-ethylpiperidino)-3-{[(4-fluorobenzyl)amino]
    methyl}benzenesulfonamide maleate. mp. 140-142°C
    N-(3,3-dimethylpiperidino)-3-{[(cyclohexylmethyl)
    amino]methyl}benzenesulfonamide hydrochloride. mp. 147-
15
    149°C
    N-(3-methylpiperidino)-4-{[(4-chlorophenethyl)amino]
    methyl}benzenesulfonamide maleate. mp. 176-178^{\circ}C
    N-(3-methylpiperidino)-4-\{[(4-chlorophenethyl)-N-
    methylamino]methyl}benzenesulfonamide maleate. mp. 168-
20
    170°C
    3-{[[2-(dimethylamino)ethyl](4-fluorobenzyl)
    amino]methyl}-N-3,3-dimethylpiperidino-
    benzenesulfonamide maleate as an oil. Mass
    spectrum(MH+=462(10%)) (TSP+)
25
    3-{[[2-(dimethylamino)ethyl](cyclohexylmethyl)
```

amino]methyl}-N-3,3-dimethylpiperidino-benzenesulfonamide maleate. mp. 149-151°C

5 EXAMPLE 4

4-{[[2-(piperidino)ethyl](2-[3,4-dimethoxy]phenylethyl)amino]methyl}-N, N-di-n-propylbenzene sulfonamide dihydrochloride

10

15

To solution of N, N-di-n-propyl-4-{[(3,4-dimethoxyphenethyl)amino]methyl}benzene sulfonamide(550 mg, 1mmole) in dry acetonitrile (100ml) was added sodium carbonate (440mg, 4.4mmole), potassium iodide (166mg,

- immole) and 2-chloroethylpiperidine hydrochloride (184mg, 1mmole). The reaction was stirred and heated at reflux for 18 hours. The reaction was poured into ice water and extracted with ethyl acetate, washed with brine, dried and evaporated to dryness *in-vacuo*.
- Chromatography on flash silica by elution with 10%methanol/dichloromethane gave 4-{[[2-(piperidino)ethyl](2-[3,4-dimethoxy]phenylethyl)amino] methyl}-N,N-di-n-propylbenzenesulfonamide which was crystallised as its dihydrochloride salt. mp. 135-137°C

EXAMPLE 5

 $N-\{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl\}-N-(4-interval)$ methylbenzyl)amine

5

10

15

Α mixture of 0.15 a М solution of 3 - [(3, 3 dimethylpiperidin-1-yl)sulfonyl]benzaldehyde in methanol (0.25 ml) and a 0.1 M solution of 4-methylbenzylamine in methanol (0.25 ml) was stirred at room temperature for 1A 0.15 M solution of sodium borohydride in methanol (0.25 ml) was added and stirring continued for a further 16 hours. The mixture was then applied to a methanol-washed 500 mg SCX solid phase extraction (SPE) cartridge and washed through with methanol (2.5 ml). The cartridge was then eluted with a 2 M solution of ammonia in methanol (2.5 ml). The eluate was vacuum evaporated to give the required product. (TS-MS: m/z $387, [M+H]^+).$

20

The following compounds were similarly prepared (mass spectrum values are given in brackets).

 $N-\{3-[(3,3-\text{dimethylpiperidin}-1-y1) \text{ sulfonyl}\}-N-[3-y]$ (4-methylpiperazin-1-yl)propyl]amine (423)

```
N-\{3-[(3,3-\text{dimethylpiperidin}-1-y1) \text{ sulfonyl}\} - N-\{3-[(3,3-\text{dimethylpiperidin}-1-y1) \text{ sulfo
```

 $N-(4-\text{chlorobenzyl})-N-\{3-[(3,3-\text{dimethylpiperidin-1-}$ 5 yl)sulfonyl]benzyl}amine (407/408)

 $N-(\text{cyclohexylmethyl})-N-\{3-[(3,3-\text{dimethylpiperidin-1-yl})\text{sulfonyl}\}$ amine (379)

10 $N=\{3-[(3,3-\text{dimethylpiperidin}-1-y1) \text{ sulfonyl}\}-N=[3-(1H-\text{imidazol}-1-y1) \text{ propyl}]$ amine (391)

N-butyl-N-{3-[(3,3-dimethylpiperidin-1yl)sulfonyl]benzyl}amine (339)

N-(tert-butyl)-N-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}amine (339)

 $N-(2-\text{chlorobenzyl})-N-\{3-[(3,3-\text{dimethylpiperidin}-1-20 yl)sulfonyl]benzyl\}amine (407/408)$

 $N-(4-\text{chlorophenethyl})-N-\{3-[(3,3-\text{dimethylpiperidin-1-yl})\ \text{sulfonyl}\}\ \text{amine}\ (421/422)$

WO 01/04087 PCT/GB00/02361

 $N-(2-\text{chlorophenethyl})-N-\{3-[(3,3-\text{dimethylpiperidin-1-yl}) \text{ sulfonyl}\}$ amine (421/422)

 $N-(2,4-\text{dichlorobenzyl})-N-\{3-[(3,3-\text{dimethylpiperidin-1-}]\}$ yl)sulfonyl]benzyl}amine (442)

 $N-\{3-[(3,3-\text{dimethylpiperidin}-1-yl) \text{ sulfonyl}\}-N-isopentylamine (353)$

10 $N-\{3-[(3,3-\text{dimethylpiperidin}-1-y1) \text{ sulfonyl}\}-N-(3-\text{methoxypropyl})$ amine (355)

 $N-\{3-[(3,3-\text{dimethylpiperidin}-1-yl) \text{ sulfonyl}\}-N-(2-\text{methylbenzyl})$ amine (387)

 $N-\{3-[(3,3-\text{dimethylpiperidin}-1-y1)\text{ sulfonyl}\}-N-(3-\text{methylcyclohexyl})$ amine (379)

 $N-\{3-[(3,3-\text{dimethylpiperidin}-1-yl) \text{ sulfonyl}\}-N-20$ hexylamine (367)

 $N-\{3-[(3,3-\text{dimethylpiperidin}-1-yl) \text{ sulfonyl}\}-N-$ propylamine (325)

N- $\{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl\}-N-(4-methylphenethyl)amine (401)$

 $N-\{3-[(3,3-\text{dimethylpiperidin}-1-y1) \text{ sulfonyl}] \text{ benzyl}\}-N-[3-(\text{trifluoromethyl}) \text{ benzyl}] \text{ amine } (441)$

5 $N-\{3-[(3,3-\text{dimethylpiperidin}-1-yl) \text{ sulfonyl}] \text{ benzyl}\}-N-[3-(\text{trifluoromethyl}) \text{ phenethyl}] \text{ amine } (455)$

EXAMPLE 6

mixture

of

а

Α

10 1-({3-[(4-benzylpiperidin-1-yl)methyl]phenyl}sulfonyl)3,3-dimethylpiperidine

0.15

M

solution

of

3 - [(3, 3 -

dimethylpiperidin-1-yl)sulfonyl]benzaldehyde in dichloromethane (0.25 ml), a 0.1 M solution of 4benzylpiperidine in dichloromethane (0.25 ml) and a 0.15 Μ solution of sodium tri-acetoxyborohydride dichloromethane (0.25 ml) was stirred at temperature for 22 hours. Methanol (1 ml) was added and the mixture applied to a methanol-washed 500 mg SCX 20 solid phase extraction (SPE) cartridge and washed through with methanol (2.5 ml). The cartridge was then eluted with a 2 M solution of ammonia in methanol (2.5 ml). The eluate was vacuum evaporated to give the 25 required product. (TS-MS: m/z 441, [M+H]).

The following compounds were similarly prepared (mass spectrum values are given in brackets).

- 2-(butyl{3-[(3,3-dimethylpiperidin-1-
- 5 yl)sulfonyl]benzyl}amino)ethan-1-ol (383)
 - 2-(benzyl{3-[(3,3-dimethylpiperidin-1-
 - yl)sulfonyl]benzyl}amino)ethan-1-ol (417)
- 10 $N-\{3-[(3,3-\text{dimethylpiperidin}-1-yl) \text{ sulfonyl}] \text{ benzyl}\}-N, N-$ bis (2-methoxyethyl) amine (399)
 - 1-(3,4-dichlorophenyl)-4-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}piperazine (497)
- $N-\{3-[(3,3-\text{dimethylpiperidin}-1-yl) \text{ sulfonyl}\}-N-$ ethyl-N-(pyridin-4-ylmethyl) amine (402)
- 1-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}-4-(4-
- 20 fluorophenyl)piperazine (446)
 - 4-{3-[(3,3-dimethylpiperidin-1-
 - yl)sulfonyl]benzyl}piperazine-1-carbaldehyde (380)
- 25 4-{3-[(3,3-dimethylpiperidin-1
 - yl)sulfonyl]benzyl}morpholine (353)

```
1-[4-(4-{3-[(3,3-dimethylpiperidin-1-
        yl)sulfonyl]benzyl)piperazin-1-yl)phenyl]ethan-1-one
         (470)
 5
         3,3-dimethyl-1-{[3-(pyrrolidin-1-
         ylmethyl)phenyl]sulfonyl)piperidine (337)
         2-{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl}-
10
         1,2,3,4-tetrahydroisoquinoline (399)
         N-\{3-[(3,3-\text{dimethylpiperidin}-1-y1) \text{ sulfonyl}\} - N, N-
         dipropylamine (367)
15
         1-benzhydryl-4-{3-[(3,3-dimethylpiperidin-1-
         yl)sulfonyl]benzyl}piperazine (518)
         N-\{3-[(3,3-dimethylpiperidin-1-yl)sulfonyl]benzyl\}-N-(2-yl)sulfonyl]benzyl\}-N-(2-yl)sulfonyl]benzyl}-N-(2-yl)sulfonyl]benzyl}-N-(2-yl)sulfonyl]benzyl}-N-(2-yl)sulfonyl]benzyl}-N-(2-yl)sulfonyl]benzyl}-N-(2-yl)sulfonyl]benzyl}-N-(2-yl)sulfonyl]benzyl}-N-(2-yl)sulfonyl]benzyl}-N-(2-yl)sulfonyl]benzyl}-N-(2-yl)sulfonyl]benzyl}-N-(2-yl)sulfonyl]benzyl}-N-(2-yl)sulfonyl]benzyl}-N-(2-yl)sulfonyl]benzyl}-N-(2-yl)sulfonyl]benzyl}-N-(2-yl)sulfonyl]benzyl}-N-(2-yl)sulfonyl]benzyl}-N-(2-yl)sulfonyl]benzyl}-N-(2-yl)sulfonyl]benzyl}-N-(2-yl)sulfonyl]benzyl}-N-(2-yl)sulfonyl]benzyl}-N-(2-yl)sulfonyl]benzyl}-N-(2-yl)sulfonyl]benzyl}-N-(2-yl)sulfonyl]benzyl}-N-(2-yl)sulfonyl]benzyl}-N-(2-yl)sulfonyl]benzyl
         methoxyethyl) - N-propylamine (383)
20
         EXAMPLE 7
         1-{3-[(3,3-dimethylpiperidin-1-
         yl) sulfonyl]benzyl}piperidine-4-carboxamide
```

10

Α mixture of a 0.15 M solution of 3-[(3,3dimethylpiperidin-1-yl)sulfonyl]benzaldehyde in methanol (0.25 ml), a 0.1 M solution of piperidine-4-carboxamide in methanol/acetic acid 4:1 v/v (0.25 ml) and a 0.15 M solution of sodium cyanoborohydride in methanol (0.25 ml) was stirred at room temperature for 18 hours. The mixture applied to a methanol-washed 500 mg SCX solid phase extraction (SPE) cartridge and washed through with methanol (2.5 ml). The cartridge was then eluted with a 2 M solution of ammonia in methanol (2.5 ml) and the eluate vacuum evaporated. The residue was dissolved in chloroform (2 ml) and the solution added to isocyanatomethyl-polystyrene (loading 1 mmole/g, mg). The suspension was shaken at room temperature for 16 hours, then filtered. The resin was washed with chloroform (2 x 2 ml) and the combined filtrates vacuum evaporated to give the required product. (TS-MS: m/z $394, [M+H]^+).$

20

The following Examples illustrate typical formulations containing a compound of the invention.

25 EXAMPLE 8

Tablets each containing 10 mg of active ingredient are made up as follows:

	Active ingredient	10 mg
5	Starch	160 mg
	Microcrystalline cellulose	100 mg
	Polyvinylpyrrolidone (as 10% solution in water)	13 mg
	Sodium carboxymethyl starch	14 mg
	Magnesium stearate	3 mg
.0		
	Total	300 mg

The active ingredient, starch and cellulose are mixed

thoroughly. The solution of polyvinylpyrrolidone is
mixed with the resultant powders and passed through a
sieve. The granules so produced are dried and re-passed
through a sieve. The sodium carboxymethyl starch and
magnesium stearate are then added to the granules which,

after mixing, are compressed on a tablet machine to
yield tablets each weighing 300 mg.

EXAMPLE 9

25 Capsules each containing 20 mg of active ingredient are made as follows:

	Active ingredient	20 mg
	Dried starch	178 mg
	Magnesium stearate	2 mg
5		
	Total	200 mg

The active ingredient, starch and magnesium stearate are

10 passed through a sieve and filled into hard gelatine
capsules in 200 mg quantities.

EXAMPLE 10

15

Capsules each containing 20 mg of medicament are made as follows:

	Active ingredient	20	mg
20	Lactose	171	mg
	Sodium lauryl sulphate	2	mg
	Sodium starch glycollate	6	mg
	Magnesium stearate	1	mg
25		200	mg

WO 01/04087

25

The active ingredient, lactose, sodium lauryl sulphate and sodium starch glycollate are mixed thoroughly. The blend is mixed with the magnesium stearate and filled into hard gelatine capsules in 200 mg quantities.

EXAMPLE 11

Tablets each containing 20 mg and medicaments are made 10 as follows:

	Active ingredient	20 mg
	Lactose	103 mg
	Microcrystalline cellulose	150 mg
15	Hydroxypropylmethylcellulose	15 mg
	Sodium starch glycollate	9 mg
	Magnesium stearate	3 mg
		300 mg
20		

The active ingredient, lactose, microcrystalline cellulose, sodium starch glycollate and hydroxypropylmethylcellulose are passed through a sieve and blended together. Water is added to the blended powders to form a damp mass. The damp mass is passed

through a coarse screen, dried, then re-screened. The dried granules are mixed with the magnesium stearate and compressed into tablets of 300 mg weight.

PCT/GB00/02361

CLAIMS

1. A compound of the formula

in which the aminosulfonyl group is attached at the 3- or 4-position, and in which

10 R^1 is hydrogen, C_{1-6} alkyl, C_{3-10} cycloalkyl, C_{3-10} cycloalkyl- C_{1-4} alkyl or optionally substituted phenyl- C_{1-4} alkyl,

 $\rm R^2$ is $\rm C_{1-6}$ alkyl, $\rm C_{3-10}$ cycloalkyl, $\rm C_{3-10}$ cycloalkyl- $\rm C_{1-4}$ alkyl, optionally substituted

5

15

15

phenyl-C₁₋₄ alkyl or -(CH₂)₂NR⁵R⁶ where R⁵ and R⁶ are each hydrogen or C₁₋₆ alkyl, and

 R^3 and R^4 are each C_{1-6} alkyl, C_{3-10} cycloalkyl, C_{3-10} cycloalkyl- C_{1-4} alkyl, C_{3-6} alkenyl, optionally substituted phenyl or optionally substituted phenyl- C_{1-4} alkyl,

or R¹ and R², or R³ and R⁴, or R⁵ and R⁶, together with the nitrogen atom to which they are attached, form a carbocyclic group containing 4 to 7 carbon atoms optionally substituted with one to three methyl or ethyl groups and optionally containing an oxygen atom or a further nitrogen atom, said carbocyclic group being optionally fused to an optionally substituted phenyl group;

or a salt thereof.

20 2. A compound according to Claim 1 in which R^1 , R^2 , R^3 and R^4 are each C_{1-6} alkyl, C_{3-10} cycloalkyl, C_{3-10} cycloalkyl- C_{1-4} alkyl or optionally substituted phenyl- C_{1-4} alkyl, and R^1 can in addition be hydrogen, or R^1 and R^2 , or R^3 and R^4 together with

the nitrogen atom to which they are attached, form a carbocyclic group.

- 3. A compound according to Claim 2 in which R^1 , R^2 , R^3 and R^4 are each C_{1-6} alkyl, C_{3-10} cycloalkyl- C_{1-4} alkyl or optionally substituted phenyl- C_{1-4} alkyl, and R^1 can in addition be hydrogen.
- 10 4. A compound according to Claim 3 in which R^1 is hydrogen, R^2 is optionally substituted phenyl- C_{1-4} alkyl and R^3 and R^4 are C_{1-6} alkyl.
 - 5. A compound according to Claim 1 in which R^2 is $-(CH_2)_2NR^5R^6$.
- 6. A compound according to Claim 1 or 5 in which R³ or R⁴ is C₃₋₆ alkyl or when R³ and R⁴ are taken together with the nitrogen atom they form a piperidine ring which is substituted at the 3-and/or 5-positions with one or two methyl or ethyl substituents.

7. A pharmaceutical formulation comprising a compound according to any of Claims 1 to 6 or a pharmaceutically acceptable salt thereof, together with a diluent or carrier therefor.

5

- 8. A compound according to any of Claims 1 to 6, for use as a pharmaceutical.
- Use of a compound according to any of Claims 1 to
 6, in the manufacture of a medicament for treating a disorder of the central nervous system.
 - 10. A method of treating a disorder of the central nervous system which comprises administering an effective amount of a compound according to Claim 1, or a pharmaceutically acceptable salt thereof.

+

Approved for use through 9/30/98. OMB 0651-0032
Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

		Patent and Tra	ademark Office:	U.S. DEF	PARTMENT OF	COMMENCE			
	A	Attorney Docket N	lumber	G-1344					
DECLARATION FOR	} <u>F</u>	irst Named Inver	ntor	Milutinovic, et al.					
UTILITY OR DESIGN	J	COMPLETE IF KNOWN							
PATENT APPLICATION)N	Application Numb	er						
	F	iling Date							
Declaration Submitted with Initial Filing		Group Art Unit							
Declaration Submitted after Initial Filing	E	Examiner Name							
As a below named inventor, I hereby declare th	nat:								
My residence, post office address, and citizenship	are as stated belo	ow next to my name.							
I believe I am the original, first and sole Inventor (i below) of the subject matter which is claimed and	if only one name is for which a patent	s listed below) or an or is sought on the Inver	iginal, first and j ntion entitled:	joint invent	tor (if plural nam	es are listed			
SULFONAMIDE SUBSTITU	TED BENZYLAMI	NE DERIVATIVES AN	D THEIR USE	AS MEDIC	CAMENTS				
the specification of which is attached hereto OR Was filed on (MM/DD/YYYY) Application Number PCT/GB00/02361 and was amended on (MM/DD/YYYY) If hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment specifically referred to above. I advinowledge the duty to disclose information which is material to patentability as defined in Title 37 Code of Federal Regulations, § 1.56.									
I hereby claim foreign priority benefits under Title Inventor's certificate, or § 365(a) of any PCT inter America, listed below and have also identified bel PCT international application having a filing date	national applications, by checking t	on which designated at ne box, any foreign ap	least one count olication for pate	try other th ent or invei	ian the United St	tates of			
	intry	Foreign Filing D	ate Pri	iority Claimed	Certified Co	py Attached NO			
Till te									
						<u> </u>			
Additional foreign application numbers are									
I hereby claim the benefit under Title 35, United S	States Code § 119	(e) of any United State	es provisional ar	oplications	(s) listed below.				
Application Number(s)	Filing D	ate (MM/DD/YYYY)		l Autori		. liontina			
9916434.5		07/13/1999	olication supplemental ereto.						

Approved for use through 9/30/98. OMB 0651-0032 Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

DECLARATION

I hereby claim the benefit under Title 35, United States Code §120 of any United States application(s), or § 365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of Title 35, United States Code § 112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations § 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. Parent Application Number	PCT Parent Number	Parent Filing Date (MM/DD/YYYY)	Parent Patent Number (if applicable)

Additional U.S. or PCT international application numbers are listed on a supplemental priority sheet attached hereto.

As a named inventor, I hereby appoint the following registered practitioner(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

Attorney Name	Reg. No.
Arvie J. Anderson	45,263
Lynn D. Apelgren	45,341
Robert A. Armitage	27,417
Brian P. Barrett	39,597
Michael T. Bates	34 ,121
Roger S. Benjamin	27,025_
Gary M. Birch	<u>48,881</u>
William R. Boudreaux	35,796
Steven P. Caltrider	3 6,467 -
Paul R. Cantrell	36,470
Charles E. Cohen	34,565
Donald L. Corneglio	_30,741
Gregory A. Cox	47,504
Paula K. Davis	47,517
Elizabeth A. Dawalt	44,646
John C. Demeter	30,167
Manisha A. Desai	43,585
Joanne Longo Feeney	_35,134
Paul J. Gaylo	_36,808_
Francis O. Ginah	44,712_
Janet A. Gongola	48,436
Arny E. Hamilton	.33,894
Frederick D. Hunter	26,915
Thomas E. Jackson	_33,064
Charles Joyner	_30,466
Gerald P. Keleher	4 <u>3,707</u>
Control of the Contro	
incomplete	

Attorney Name	Reg. No.
James J. Kelley	41,888
Paul J. Koivuniemi	_31,533_
Robert E. Lee	_27,919_
Kirby Lee	<u>47,744</u>
James P. Leeds	35,241
Nelsen L. Lentz	38,537_
Douglas K. Norman	33,267
Arleen Palmberg	40,422
Thomas G. Plant	35,784
Edward Prein	37,212
Grant E. Reed	41,264
James J. Sales	33,773
Michael J. Sayles	32,295
Robert L. Sharp	45,609
David M. Stemerick	40,187
Mark J. Stewart	43,936
Robert D. Titus	_40,206_
Robert C. Tucker	-45 ,16 5
Tina M. Tucker	-47 <u>.145</u>
MaCharri Vorndran-Jones	_36,711_
Gilbert T. Voy	43,972
Thomas D. Webster	39.872
Lawrence T. Welch	29,487
Alexander Wilson	45,782
Dan L. Wood	48,613

Direct al	l corresponde	nce to:								
Name) ,	ELILILLY AND	COMPANY	2						
Addre	ess	ATTN: Janet A.	Gongola							
Addre	ess	LILLY CORPOR	ATE CENTE	R/DC1104						
City		INDIANAPOLIS	St	State INDIANA				4	<u>16285</u>	
Coun		US at all statements made	Telep			(317) 277-6		Fax	(317) 276	
jeoparc	dize the valid	or imprisonment, or bo dity of the application of or First Inventor:	or any patent iss						miu iaise statei	nente may
					de been me	a ioi tilis alisi	g	Cilcoi		
Given Name	Sar	ndra		Middle Name	Ginette		Milutin		Suffix e.g. Jr.	
Name	Sar or's Signat	ndra	Cerpner.			Family	-			2-2001
Name Invent	-	ndra	Cerhner.	Name		Family	Milutin	ovic	e.g. Jr.	2 200\ FR
Name Invent	or's Signat	ure 37	Cerpner .	Name	Ginette	Family Name	Milutin	ovic	e.g. Jr. 3 – №	
Name Invento Reside Addres	or's Signat	Reading 39 Devonsh	Cennai (s s	Name	Ginette	Family Name	Milutin	ovic	e.g. Jr. 3 – №	

O

Approved for use through 9/30/98. OMB 0651-0032 Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

DECLARATION																
Name of Additional Joint Inventor, if any:								A Petition has been filed for this unsigned inventor							ventor	
Given Robin							acorde John House									
r's re		(2)	Til m	b									Date		<u> </u>	
nce: City	٧	Voking	ham			Sta	ate	Ber	kshire	. Cc	ountry	UK C	13X	Citize	enship	UK
Address 2 Plover Close,																
fice Addr	ess	SAME	AS /	ABO	VΕ											
Wokin	gha	am		State	Berk	shi	ire		Postal C	stal Code RG41 3JD		Country	UK			
of Addi	tior	nal Join	Inve	entor	, if any	7:				A Pet	tition ha	s been f	iled for thi			ventor
Da	vid								Edwar	u I	Family Name	Tuppe	<u>er</u>			
r's ire		an			-Ca	,	, F		4-				Date	3,3	1110	7
	R	eading		i		T -	erks	hire		Cour	ntry Uk	(Cel	3+	Citiz	enship	UK
ffice Addr	ess	8 Mild	lenha	all C	lose,	Lov	ver	Earle	; у,							
ffice Addr	ess	SAME	AS	ABO	VE											
Readi	ng		State	E	Berksh	ire			Postal (Code	RG6	зат	Country	, U	K	
	Rolling Rollin	Robin r's re dice: City V s fice Address Wokingha of Addition David	Robin T's Vocation of Ploy Flice Address SAME Wokingham Of Additional Joint David T's Joint Playid Tress The Ploy Reading Flice Address SAME Reading Flice Address SAME	Robin T's Vokingham 2 Plover C Fice Address SAME AS Wokingham of Additional Joint Invented Playing Ince: City Reading Fice Address SAME AS Fifice Address SAME AS	Robin T's Vokingham 2 Plover Close Flice Address SAME AS ABO Wokingham State of Additional Joint Inventor David T's Ince: City Reading Flice Address SAME AS ABO Flice Address SAME AS ABO	Robin T's Core City Wokingham 2 Plover Close, Flice Address SAME AS ABOVE Wokingham State Berk of Additional Joint Inventor, if any David T's Free City Reading State Flice Address SAME AS ABOVE Flice Address SAME AS ABOVE	Robin T's Wokingham State 2 Plover Close, SAME AS ABOVE Wokingham State Berkshi of Additional Joint Inventor, if any: David T's Tree Total Control Close, Berkshi of Additional Joint Inventor, if any: David T's Tree Total Control Close, SAME AS ABOVE Fiice Address SAME AS ABOVE	of Additional Joint Inventor, if any: Robin	of Additional Joint Inventor, if any: Robin	Robin Middle Rame George re	Robin Middle Name George T's Robin State Berkshire Countries SAME AS ABOVE Wokingham State Berkshire Postal Code of Additional Joint Inventor, if any: A Period Additional Joint Inventor, if any:	of Additional Joint Inventor, if any: Robin	of Additional Joint Inventor, if any: Robin	A Petition has been filed for this Robin	A Petition has been filed for this unsigned for the sunsigned for	A Petition has been filed for this unsigned in Middle Name George Family Name Simmonds e.g. Jr. Robin